



7322 Highway 1
Coldbrook, NS B4R 1B9

Dr. Scott Schofield, BScH, DDS
Dr. Elizabeth Jackson, BScH, DDS
Dr. April Nason, BSc, DDS

Extractions – Consent

I have recommended one or more of your teeth be extracted based upon your symptoms, my examination of your mouth, the treatment plan I have discussed with you, and your choice. Uncomplicated extractions are a predictable minor surgery. I, nevertheless, want you to be aware of the commonly known risks and side effects of this procedure. On occasion a tooth requires surgical removal where an incision is made in gum tissue or bone is removed to gain access to the tooth.

You may experience **pain, swelling** or **bleeding** for a time after the extraction. Instructions on how to manage these problems, if they occur, are described on the back of this page. These should last for only a short while, but should a problem become more severe or last longer than you anticipated, call our office immediately.

You may experience an **infection** following the extraction. I will advise you what to look for as a sign of infection. If any of these signs occur you should call or see me as soon as possible. Infections can involve aggressive bacteria and in some cases may require hospitalization to successfully treat.

Teeth adjacent to the extraction tooth may be chipped, damaged or lost during the extractions.

Nerves which supply sensation to your mouth, chin, lips, tongue, and gum tissue may run near the area of extraction. After the extraction you may experience **alteration of normal sensation** (itching, burning or tingling, for example) for a short or indefinite period of time. In very rare cases the sensation may be lost permanently.

You may experience a painful but harmless condition known as a **dry socket**. This occurs when the protective blood clot is dislodged, exposing and irritating nerve endings. Closely following the post-operative instructions minimizes the risk of developing this condition. Although dry sockets are temporary and not harmful, it is painful. It can be readily treated and you should seek treatment from me. I will place a medicine in the socket that will soothe and protect it while alleviating the pain.

For teeth in the upper jaw there is a risk that following the extraction a hole or pathway may be present between the sinus and your oral cavity. This is because the roots of upper teeth end just below the floor of the sinus and sometimes actually go through the sinus floor. If a **sinus perforation** does occur, I may need to make a small surgical repair of the hole and may place you on antibiotics and antihistamines to reduce the risk of a sinus infection.

Following the procedure the muscles of your **jaw may be stiff or sore** and it may be difficult to open your mouth wide for several days. This is a temporary condition and moist heat and pain medication may provide some relief. You may also experience some cracking or redness in the corners of your mouth.

I invite your questions concerning the risks discussed and contained in this document. By signing below you acknowledge that you have read and understand the information, and have had all your questions answered satisfactorily.

Additional comments: _____

Date: _____

Patient: _____

Dentist: _____